

Publicacions Matemàtiques, Vol **41** (1997), 519–526.

THE VORONOVSKAYA THEOREM FOR SOME LINEAR POSITIVE OPERATORS IN EXPONENTIAL WEIGHT SPACES

L. REMPULSKA AND M. SKORUPKA

Abstract

In this note we give the Voronovskaya theorem for some linear positive operators of the Szasz-Mirakjan type defined in the space of functions continuous on $[0, +\infty)$ and having the exponential growth at infinity.

Some approximation properties of these operators are given in [3], [4].

1. Preliminaries

1.1. Let $R_0 := [0, +\infty)$, $N := \{1, 2, \dots\}$, $N_0 := N \cup \{0\}$ and let $w_r(\cdot)$, $r > 0$, be the weight function defined on R_0 by the formula

$$(1) \quad w_r(x) := e^{-rx}.$$

Similarly as in [1] we denote by C_r , $r > 0$, the space of real-valued functions f defined on R_0 and such that $w_r f$ is a uniformly continuous and bounded function on R_0 . The norm in C_r is defined by

$$\|f\|_r := \sup_{x \in R_0} w_r(x) |f(x)|.$$

For a fixed $r > 0$ let

$$C_r^2 := \{f \in C_r : f', f'' \in C_r\}.$$

1.2. In [3] were introduced the following operators of the Szasz-Mirakjan type for functions $f \in C_r$, $r > 0$,

$$(2) \quad L_n^{(1)}(f; x) := \sum_{k=0}^{\infty} p_{n,k}(x) f\left(\frac{2k}{n}\right),$$

$$(3) \quad L_n^{(2)}(f; x) := \sum_{k=0}^{\infty} p_{n,k}(x) \frac{n}{2} \int_{I_{n,k}} f(t) dt,$$

Keywords. Voronovskaya theorem, linear positive operator.

1991 *Mathematics subject classifications*: 41A36.

$x \in R_0$, $n \in N$, where

$$(4) \quad p_{n,k}(x) := \frac{1}{\cosh nx} \frac{(nx)^{2k}}{(2k)!}, \quad k \in N_0,$$

$\sinh x$, $\cosh x$, $\tanh x$ are the elementary hyperbolic functions and $I_{n,k} := \left[\frac{2k}{n}, \frac{2k+2}{n}\right]$, $k \in N_0$.

In [4] were introduced the operators

$$(5) \quad L_n^{(3)}(f; x) := \frac{f(0)}{1 + \sinh nx} + \sum_{k=0}^{\infty} q_{n,k}(x) f\left(\frac{2k+1}{n}\right),$$

$$(6) \quad L_n^{(4)}(f; x) := \frac{f(0)}{1 + \sinh nx} + \sum_{k=0}^{\infty} q_{n,k}(x) \frac{n}{2} \int_{I_{n,k}^*} f(t) dt,$$

$x \in R_0$, $n \in N$, where

$$(7) \quad q_{n,k}(x) := \frac{1}{1 + \sinh nx} \frac{(nx)^{2k+1}}{(2k+1)!},$$

and $I_{n,k}^* := \left[\frac{2k+1}{n}, \frac{2k+3}{n}\right]$ for $k \in N_0$.

We observe that the above operators are linear positive operators well-defined on every space C_r , $r > 0$, and

$$(8) \quad L_n^{(i)}(1; x) = 1, \quad 1 \leq i \leq 4,$$

for all $x \in R_0$ and $n \in N$.

In [3] and [4] it was proved that $L_n^{(i)}$, $1 \leq i \leq 4$, are operators from C_r into C_s for every fixed $s > r > 0$ provided n is large enough. Moreover in [3], [4] some approximation properties of these operators were given. In particular in [3], [4] we proved the following

Theorem A. *Suppose that r, s, n_0 are fixed numbers such that $s > r > 0$, $n_0 \in N$ and $n_0 > r \left(\ln \frac{s}{r}\right)^{-1}$. If $f \in C_r$, then there exists a positive constant $M_1 \equiv M_1(n_0, r, s)$ depending only on n_0, r, s such that for all $x \in R_0$, $n_0 < n \in N$ and $1 \leq i \leq 4$*

$$w_s(x) \left| L_n^{(i)}(f; x) - f(x) \right| \leq M_1 \omega \left(f, C_r; \sqrt{\frac{x+1}{n}} \right),$$

where $\omega(f; C_r; \cdot)$ is the modulus of continuity of f , i.e.,

$$\omega(f; C_r; t) := \sup_{0 < h \leq t} \|f(\cdot + h) - f(\cdot)\|_r.$$

2. Auxiliary results

In this part we shall give some properties of the operators $L_n^{(i)}$. Let

$$(9) \quad \begin{aligned} S(nx) &:= \frac{\sinh nx}{1 + \sinh nx}, \\ T(nx) &:= \frac{\cosh nx}{1 + \sinh nx}, \\ V(nx) &:= 1 - \tanh nx, \end{aligned}$$

for $n \in N$ and $x \in R_0$. By elementary calculations from (2)-(8) and (9) we obtain

Lemma 1. *For all $x \in R_0$ and $n \in N$ we have*

$$\begin{aligned} L_n^{(1)}(t-x; x) &= -xV(nx), \\ L_n^{(1)}((t-x)^2; x) &= \left(2x^2 - \frac{x}{n}\right) V(nx) + \frac{x}{n}, \\ L_n^{(1)}((t-x)^4; x) &= \left(8x^4 - \frac{12x^3}{n} + \frac{4x^2}{n^2} - \frac{x}{n^3}\right) V(nx) + \frac{3x^2}{n^2} + \frac{x}{n^3}, \\ L_n^{(2)}(t-x; x) &= -xV(nx) + \frac{1}{n}, \\ L_n^{(2)}((t-x)^2; x) &= \left(2x^2 - \frac{3x}{n}\right) V(nx) + \frac{x}{n} + \frac{4}{3n^2}, \\ L_n^{(2)}((t-x)^4; x) &= \left(8x^4 - \frac{28x^3}{n} + \frac{32x^2}{n^2} - \frac{21x}{n^3}\right) V(nx) + \frac{12x}{n^3} + \frac{16}{5n^4}, \\ L_n^{(3)}(t-x; x) &= x(T(nx) - 1), \\ L_n^{(3)}((t-x)^2; x) &= x^2(S(nx) - 2T(nx) + 1) + \frac{x}{n}V(nx), \\ L_n^{(3)}((t-x)^4; x) &= x^4(7S(nx) - 8T(nx) + 1) + \frac{12x^3}{n}(T(nx) - S(nx)) \\ &\quad + \frac{x^2}{n^2}(7S(nx) - 4T(nx)) + \frac{x}{n^3}T(nx), \\ L_n^{(4)}(t-x; x) &= x(T(nx) - 1) + \frac{1}{n}S(nx), \\ L_n^{(4)}((t-x)^2; x) &= x^2(S(nx) - 2T(nx) + 1) \\ &\quad + \frac{2x}{n}(T(nx) - S(nx)) + \frac{4}{3n^2}S(nx), \\ L_n^{(4)}((t-x)^4; x) &= x^4(7S(nx) - 8T(nx) + 1) + \frac{28x^3}{n}(T(nx) - S(nx)) \\ &\quad + \frac{x^2}{n^2}(35S(nx) - 32T(nx)) + \frac{17x}{n^3}T(nx). \end{aligned}$$

Using Lemma 1, we shall prove two lemmas.

Lemma 2. *For every fixed $x_0 \in R_0$ one has*

$$(10) \quad \lim_{n \rightarrow \infty} nL_n^{(i)}(t - x_0; x_0) = \begin{cases} 0 & \text{if } i = 1, 3, \\ 1 & \text{if } i = 2, 4, \end{cases}$$

and

$$(11) \quad \lim_{n \rightarrow \infty} nL_n^{(i)}((t - x_0)^2; x_0) = x_0 \quad \text{for } 1 \leq i \leq 4.$$

Proof: We shall prove only (10) and (11) for $i = 3$, because the proof for $i = 1, 2, 4$ is analogous.

By Lemma 1 and (9) we have

$$\begin{aligned} nL_n^{(3)}(t - x; x) &= \frac{nx}{e^{2nx}(1 + \sinh nx)} - \frac{nx}{(1 + \sinh nx)}, \\ nL_n^{(3)}((t - x)^2; x) &= \frac{nx^2}{1 + \sinh nx} - \frac{2nx^2}{e^{nx}(1 + \sinh nx)} + \frac{x \cosh nx}{1 + \sinh nx}, \end{aligned}$$

for every $x \in R_0$ and $n \in N$, which immediately yield (10) and (11). ■

Lemma 3. *For every fixed $x_0 \in R_0$ there exists a positive constant $M_2(x_0)$, depending only on x_0 , such that for all $n \in N$*

$$(12) \quad L_n^{(i)}((t - x_0)^4; x_0) \leq M_2(x_0)n^{-2}, \quad 1 \leq i \leq 4.$$

Proof: For example we shall prove (12) for $L_n^{(1)}$. By (9) we have for $n \in N$, $p \in N$ and $x \in R_0$

$$0 \leq x^p V(nx) = \frac{2x^p}{e^{2nx} + 1} \leq 2^{1-p} p! n^{-p}.$$

Applying the above inequality to the formula given in Lemma 1, we obtain

$$L_n^{(1)}((t - x_0)^4; x_0) \leq \frac{47}{n^4} + \frac{3x_0^2}{n^2} + \frac{x_0}{n^3} \leq M_2(x_0)n^{-2},$$

for every fixed $x_0 \geq 0$ and for all $n \in N$. ■

The proof of (12) for $i = 2, 3, 4$ is similar.

In the papers [3] (for $L_n^{(i)}$, $i = 1, 2$) and [4] (for $L_n^{(i)}$, $i = 3, 4$) we proved the following two lemmas.

Lemma 4. Let $s > r > 0$ and let n_0 be a natural number such that

$$(13) \quad n_0 > r \left(\ln \frac{s}{r} \right)^{-1}.$$

Then there exists a positive constant $M_3 \equiv M_3(r, s, n_0)$ depending only on r, s, n_0 such that for all $n > n_0$ and $i = 1, 2, 3, 4$

$$\left\| L_n^{(i)} \left(\frac{1}{w_r(t)}; \cdot \right) \right\|_s \leq M_3.$$

Lemma 5. Suppose that r, s and n_0 are a numbers as in Lemma 4. Then there exists a positive constant $M_4 \equiv M_4(r, s, n_0)$ depending only on r, s, n_0 such that for all $x \geq 0, n > n_0$ and $i = 1, 2, 3, 4$

$$(14) \quad w_s(x) L_n^{(i)} \left(\frac{(t-x)^2}{w_r(t)}; x \right) \leq M_3 \frac{x+1}{n}.$$

Applying the above lemmas, we shall prove

Lemma 6. Suppose that x_0 is a fixed point on R_0 and $\varphi(\cdot; x_0)$ is a function belonging to a give space $C_r, r > 0$, such that $\lim_{t \rightarrow \infty} \varphi(t; x_0) = 0, (\lim_{t \rightarrow 0+} \varphi(t; 0) = 0)$. Then

$$(15) \quad \lim_{n \rightarrow \infty} L_n^{(i)} (\varphi(t; x_0); x_0) = 0 \quad \text{for } 1 \leq i \leq 4.$$

Proof: We shall prove (15) for $i = 1$, because the proof of (15) for $i = 2, 3, 4$ is analogous.

Choose $\varepsilon > 0$ and M_3 as in Lemma 4. Then by the properties of $\varphi(\cdot; x_0)$ there exist positive constants $\delta \equiv \delta(\varepsilon, M_3)$ and M_5 such that

$$w_r(t) |\varphi(t; x_0)| < \frac{\varepsilon}{2 M_3} \quad \text{for } |t - x_0| < \delta,$$

$$w_r(t) |\varphi(t; x_0)| < M_5 \quad \text{for } t \geq 0.$$

Denoting by $Q_{n,1} := \{k \in N_0 : |\frac{2k}{n} - x_0| < \delta\}$ and $Q_{n,2} := \{k \in N_0 : |\frac{2k}{n} - x_0| \geq \delta\}$, we get for $s > r$ and $n > n_0$ by (1)-(4) and Lemma 4

$$\begin{aligned} w_s(x_0) \left| L_n^{(1)} (\varphi(t; x_0); x_0) \right| &\leq w_s(x_0) \sum_{k=0}^{\infty} p_{n,k}(x_0) \left| \varphi \left(\frac{2k}{n}; x_0 \right) \right| \\ &= w_s(x_0) \sum_{k \in Q_{n,1}} p_{n,k}(x_0) \left| \varphi \left(\frac{2k}{n}; x_0 \right) \right| \\ &\quad + w_s(x_0) \sum_{k \in Q_{n,2}} p_{n,k}(x_0) \left| \varphi \left(\frac{2k}{n}; x_0 \right) \right| \\ &:= \sum_1 + \sum_2 \end{aligned}$$

and

$$\begin{aligned} \sum_1 &< \frac{\varepsilon}{2M_3} w_s(x_0) \sum_{k=0}^{\infty} p_{n,k}(x_0) \left(w_r \left(\frac{2k}{n} \right) \right)^{-1} < \frac{\varepsilon}{2}, \\ \sum_2 &\leq M_5 w_s(x_0) \sum_{k \in Q_{n,2}} p_{n,k}(x_0) \left(w_r \left(\frac{2k}{n} \right) \right)^{-1}. \end{aligned}$$

Since $1 \leq \delta^{-2} \left(\frac{2k}{n} - x_0 \right)^2$ if $\left| \frac{2k}{n} - x_0 \right| \geq \delta$, we have

$$\begin{aligned} \sum_2 &\leq M_5 \delta^{-2} w_s(x_0) \sum_{k \in Q_{n,2}} p_{n,k}(x_0) \left(w_r \left(\frac{2k}{n} \right) \right)^{-1} \left(\frac{2k}{n} - x_0 \right)^2 \\ &\leq M_5 \delta^{-2} w_s(x_0) L_n^{(1)} \left(\frac{(t-x_0)^2}{w_r(t)}; x_0 \right), \end{aligned}$$

wich by (14) and (13) yields

$$\sum_2 \leq M_5 M_4 \frac{x_0 + 1}{n \delta^2} \quad \text{for all } n > n_0.$$

It is obvious that for fixed numbers $\varepsilon > 0$, $\delta > 0$, $M_3 > 0$, $M_4 > 0$, $n_0 \in N$ and $x_0 \geq 0$ there exist a natural number $n_1 > n_0$ depending on the above parameters such that for all $n_1 < n \in N$

$$M_4 M_5 \frac{x_0 + 1}{n \delta^2} < \frac{\varepsilon}{2}.$$

Hence we have

$$\sum_2 < \frac{\varepsilon}{2} \quad \text{for all } n > n_1.$$

Consequently,

$$w_s(x_0) |L_n^{(1)}(\varphi(t; x_0); x_0)| < \varepsilon \quad \text{for } n > n_1,$$

which proves that

$$\lim_{n \rightarrow \infty} w_s(x_0) L_n^{(1)}(\varphi(t; x_0); x_0) = 0.$$

From this and (1) assertion (15) follows for x_0 and $i = 1$. Thus the proof is completed. ■

3. Theorems of the Voronovskaya type

The Voronovskaya theorem for the Bernstein operators is given in [2].

We shall prove a similar theorem for the operators $L_n^{(i)}$.

Theorem 1. *Let $f \in C_r^2$ with some $r > 0$. Then*

$$(16) \quad \lim_{n \rightarrow \infty} n \left\{ L_n^{(i)}(f; x) - f(x) \right\} = \frac{x}{2} f''(x)$$

for every $x \in R_0$ and $i = 1, 3$.

Proof: Let $x_0 \geq 0$ be an arbitrary fixed point and $i = 1$. By the Taylor formula we have for $t \geq 0$

$$(17) \quad f(t) = f(x_0) + f'(x_0)(t - x_0) + \frac{1}{2} f''(x_0)(t - x_0)^2 + \psi(t; x_0)(t - x_0)^2,$$

where $\psi(\cdot; x_0)$ is a function belonging to the space C_r and $\lim_{t \rightarrow x_0} \psi(t; x_0) = 0$. By (2), (8) and (17) we get

$$(18) \quad L_n^{(1)}(f(t); x_0) = f(x_0) + f'(x_0) L_n^{(1)}(t - x_0; x_0) + \frac{1}{2} f''(x_0) L_n^{(1)}((t - x_0)^2; x_0) + L_n^{(1)}(\psi(t; x_0)(t - x_0)^2; x_0)$$

for every $n \in N$. Using Lemma 2, we have

$$(19) \quad \begin{aligned} \lim_{n \rightarrow \infty} n L_n^{(1)}(t - x_0; x_0) &= 0, \\ \lim_{n \rightarrow \infty} n L_n^{(1)}((t - x_0)^2; x_0) &= x_0. \end{aligned}$$

By (2) and the Hölder inequality we have for every $n \in N$

$$(20) \quad \begin{aligned} &\left| L_n^{(1)}(\psi(t; x_0)(t - x_0)^2; x_0) \right| \\ &\leq \left\{ L_n^{(1)}(\psi^2(t; x_0); x_0) \right\}^{\frac{1}{2}} \left\{ L_n^{(1)}((t - x_0)^4; x_0) \right\}^{\frac{1}{2}}. \end{aligned}$$

Since for the function $\varphi(t; x_0) := \psi^2(t; x_0)$, $t \geq 0$, we have $\varphi(\cdot; x_0) \in C_{2r}$ and $\lim_{t \rightarrow x_0} \varphi(t; x_0) = 0$, we get by Lemma 6

$$(21) \quad \lim_{n \rightarrow \infty} L_n^{(1)}(\psi^2(t; x_0); x_0) \equiv \lim_{n \rightarrow \infty} L_n^{(1)}(\varphi(t; x_0); x_0) = 0.$$

Applying (21) and (12) to (20), we obtain

$$(22) \quad \lim_{n \rightarrow \infty} n L_n^{(1)}(\psi(t; x_0)(t - x_0)^2; x_0) = 0.$$

Now we immediately obtain (16) for a given x_0 and $i = 1$ from (18) by (19) and (22). This proves the desired assertion for $i = 1$. ■

Similarly we can prove the following

Theorem 2. *Suppose that $f \in C_r^2$ with some $r > 0$. Then*

$$\lim_{n \rightarrow \infty} n \left\{ L_n^{(i)}(f; x) - f(x) \right\} = f'(x) + \frac{x}{2} f''(x)$$

for every $x \in R_0$ and $i = 2, 4$.

References

1. M. BECKER, D. KUCHARSKI AND R. J. NESSEL, Global approximation theorems for the Szasz-Mirakjan operators in exponential weight spaces, in “*Linear Spaces and Approximation*,” Proc. Conf. Oberwolfach, 1977, Birkhäuser Verlag, Basel.
2. P. P. KOROVKIN, “*Linear operators and Approximation Theory*,” Moscow, 1959 (Russian).
3. M. LEŚNIEWICZ AND L. REMPULSKA, Approximation by some operators of the Szasz-Mirakjan type in exponential weight spaces, *Glas. Mat. Ser. III*, (in print).
4. L. REMPULSKA AND M. SKORUPKA, On approximation of functions by some operators of the Szasz-Mirakjan type, *Fasc. Math.* **26** (1996), 123–134.

Institute of Mathematics
Poznań University of Technology
Piotrowo 3A
60-965 Poznań
POLAND

Primera versió rebuda el 9 de Juliol de 1996,
darrera versió rebuda el 18 de Novembre de 1996